

## ACKNOWLEDGEMENTS

This work was supported in part by the Fifth Framework Programme of the European Commission (Contract No. QLK2-2000-01369), the Consejería de Salud de la Junta de Andalucía (Exp. 32/02) and the Thematic Network for Joint Research G03/90, Spanish Ministry of Health and Consumer Affairs.

## REFERENCES

- Jameson B. Serology of *Pneumocystis carinii*. In: Young LS, ed. *Pneumocystis carinii pneumonia*, 1st edn. New York: Dekker, 1984; 97–106.
- Peglow SL, Smulian AG, Linke MJ *et al.* Serologic response to *Pneumocystis carinii* antigens in health and disease. *J Infect Dis* 1990; **161**: 296–306.
- Smulian AG, Sullivan DW, Linke MJ *et al.* Geographic variation in the humoral response to *Pneumocystis*. *J Infect Dis* 1993; **167**: 1243–1247.
- Meuwissen JHE, Tauber I, Leeuwenberg ADEM, Beckers PJA, Sieben M. Parasitologic and serologic observations of infection with *Pneumocystis* in humans. *J Infect Dis* 1977; **136**: 43–49.
- Pifer LL, Hughes WT, Stagno S, Woods D. *Pneumocystis carinii* infection: evidence for high prevalence in normal and immunosuppressed children. *Pediatrics* 1978; **61**: 35–41.
- Lundgren B, Lebech M, Lind K, Nielsen JO, Lundgren JD. Antibody response to a major human *Pneumocystis carinii* antigen in patients without evidence of immunosuppression and in patients with suspected atypical pneumonia. *Eur J Clin Microbiol Infect Dis* 1993; **12**: 105–109.
- Wakefield AE, Stewart TJ, Moxon ER, Marsh K, Hopkin JM. Infection with *Pneumocystis carinii* is prevalent in healthy Gambian children. *Trans R Soc Trop Med Hyg* 1990; **84**: 800–802.
- Vargas SL, Hughes WT, Santolaya ME *et al.* Search for primary infection by *Pneumocystis carinii* in a cohort of normal healthy children. *Clin Infect Dis* 2001; **32**: 855–861.
- Calderón EJ, Regordan C, Medrano FJ, Ollero M, Varela JM. *Pneumocystis carinii* infection in patients with chronic bronchial disease. *Lancet* 1996; **347**: 977.
- Sing A, Roggenkamp A, Autenrieth IB, Heesemann J. *Pneumocystis carinii* carriage in immunocompetent patients with primary pulmonary disorders as detected by single or nested PCR. *J Clin Microbiol* 1999; **37**: 3409–3410.
- Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970; **227**: 680–685.
- Towbin H, Staehelin T, Gordon J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci USA* 1979; **76**: 4350–4354.
- Aliouat EM, Martinez A, Jiménez E *et al.* Development of pneumocystosis animal models: corticosteroid-treated Wistar rat; SCID mouse and nude rat. *J Eukaryot Microbiol* 1997; **44**(suppl): 41S–42S.
- Medrano FJ, Respaldiza N, Medrano A *et al.* Seroprevalence of *Pneumocystis* human infection in southern Spain. *J Eukaryot Microbiol* 2003; **50**(suppl): 649–650.
- Kohout E, Post C, Azadeh B, Dutz W, Banderizadeh B, Kadivar D. Immunoglobulin levels in infantile pneumocystosis. *J Clin Pathol* 1972; **25**: 135–140.
- Chatterton JMW, Joss AWL, Pennington TH, Do H-Y. Usefulness of rat derived antigen in the serodiagnosis of *Pneumocystis carinii* infection. *J Med Microbiol* 1999; **48**: 681–687.
- Gigliotti F. Host species-specific antigenic variation of a mannosylated surface glycoprotein of *Pneumocystis carinii*. *J Infect Dis* 1992; **165**: 329–336.
- Walzer PD, Kim CK, Cushion MT. *Pneumocystis carinii*. In: Walzer PD, Genta RM, eds. *Parasitic infections in the compromised host*, 1st edn. New York: Dekker, 1989; 83–178.
- Vargas SL, Ponce CA, Gigliotti F *et al.* Transmission of *Pneumocystis carinii* DNA from a patient with *P. carinii* pneumonia to immunocompetent contact health care workers. *J Clin Microbiol* 2000; **38**: 1536–1538.

## RESEARCH NOTE

### *Salmonella* Enteritidis in Bosnia and Herzegovina

S. Uzunovic-Kamberovic

Laboratory for Sanitary and Clinical Microbiology, Canton Institute Public Health, Zenica, Bosnia and Herzegovina

## ABSTRACT

A comprehensive retrospective analysis of human *Salmonella* Enteritidis isolates in the Zenica-Doboj Canton of Bosnia and Herzegovina was conducted by inquiry and questionnaire. In the period 1998–2000, 299 isolates of *Salmonella* spp. were recorded, of which *S. Enteritidis* accounted for 74.2%. The isolation rate of *S. Enteritidis* increased during this period, from 12.7 to 25.5 isolates/year/100 000 population. Isolates were obtained all year round, mostly from sporadic cases of infection or limited family outbreaks. Home-made food was identified as the most important source of infection, being implicated in 81% of outbreaks and 81.7% of cases of sporadic infection.

**Keywords** Bosnia and Herzegovina, food-borne infection, *Salmonella* Enteritidis

Corresponding author and reprint requests: S. Uzunovic-Kamberovic, Fra Ivana Jukica 2/4, 72000 Zenica, p. b. 62, Bosnia and Herzegovina  
E-mail: selma\_kamb@yahoo.com

**Original Submission:** 6 April 2004; **Accepted:** 17 May 2004

*Clin Microbiol Infect* 2004; 10: 1031–1033  
10.1111/j.1469-0691.2004.00988.x

*Salmonella* Enteritidis has emerged worldwide since 1980 as the commonest serotype isolated from cases of human food-borne disease [1,2]. There is no national *Salmonella* centre in Bosnia and Herzegovina, so surveillance of *Salmonella* infections is primarily the responsibility of the Canton Institutes for Public Health. These centres receive mandatory daily morbidity reports from general practitioners, as well as notification records of patients with *Salmonella* infection. Epidemiologists then conduct local surveys and interview members of the patient's household and other individuals who may have been exposed. A standard questionnaire is used that requests demographic and socio-economic data and records information on date of onset of illness, date of inquiry, dietary habits, food preferences, ownership of pets and other animals, consumption of food and water, and travel abroad in the 3 days before the onset of illness. In particular, patients were asked to report their own belief concerning the source of the infection.

A comprehensive retrospective analysis of all human *S. Enteritidis* isolates was conducted by inquiry and systematic examination of the questionnaires. Recorded cases were classified as either: (1) *verified*, if the agent was isolated from both an ill patient and a food source, or if the agent was isolated from either the food or the patient and a link could be determined on the basis of the information in the questionnaire; or (2) *probable*, if the association, based solely on information collected during the interview, was sufficiently strong, e.g., several cases with similar symptoms and only one meal in common.

Faecal specimens were streaked onto *Salmonella*-Shigella (SS) agar (Oxoid, Basingstoke, UK) directly and after overnight incubation at 37°C in selenite enrichment broth (Oxoid). Microbiological analysis of food products was performed routinely at the Laboratory for Sanitary and Clinical Microbiology (Canton Institute Public Health, Zenica), according to the standards and legal regulations of the Republic/Federation of Bosnia and Herzegovina. *Salmonella* was detected

in food products by homogenising 25 g of food sample in 250 mL selenite broth and incubating overnight at 37°C, and then subculturing on SS agar. Following a further incubation period overnight at 37°C, colonies were identified and serotyped by conventional microbiological methods [3]. Each patient was registered only once, even if more than one episode of infection had occurred or the patient had been registered in more than one database.

Two microbiology laboratories in the Zenica-Doboj Canton serve a total population of 331 229, divided into an urban area with 149 053 inhabitants, and a rural area with 182 176 inhabitants. In the period 1998–2000, 299 isolates of *Salmonella* spp. were recorded, of which *S. Enteritidis* accounted for 74.2%. The isolation rate of *S. Enteritidis* increased during this period from 12.7 to 25.5/year/100 000 population. The average annual age-specific rates of *S. Enteritidis* were highest among children aged <6 years in both the urban and rural areas. The cases occurred all year round, although the maximum seasonal peak was observed during September and October.

Not all infected individuals develop symptoms severe enough to need medical attention, but even when this is the case, the physician may not send the patient's stool for analysis. Thus, as many cases of *Salmonella* infection are not reported, the figures recorded (Table 1) probably represent only a fraction of the actual number of cases that occur, as has been shown in many other countries [1,4]. Most infections in this region with *S. Enteritidis* occur as sporadic cases or limited family outbreaks (mean of three cases/outbreak) rather than as part of large common-source outbreaks. However, apparently sporadic cases

**Table 1.** *Salmonella* Enteritidis infections in the Zenica-Doboj Canton, Bosnia and Herzegovina

Year	Verified no. of outbreaks	Verified no. of individuals affected in outbreaks	Probable no. of individuals affected in outbreaks <sup>a</sup> (no. of outbreaks involved)	Verified no. of sporadic cases	Probable no. of sporadic cases <sup>a</sup> (probable no. of outbreaks)
1998	5	24	1 (1)	31	10 (5)
1999	10	26	7 (3)	47	22 (11)
2000	12	24	4 (3)	86	58 (19)
1998–2000 total	27	74	11 (7)	164	90 (35)

<sup>a</sup>Individuals who did not visit a physician. Data extracted from the patient's records and from questionnaires.

may frequently represent unrecognised outbreaks (Table 1).

Home-made food was the most important source of *S. Enteritidis* infection in this region, being implicated in 22 (81%) outbreaks and 134 (81.7%) sporadic infections. This rate is higher than that in some other reports [2,5]. If imperfect practice in kitchens has contributed to the striking increase in *S. Enteritidis* food poisoning in Bosnia and Herzegovina, this may be evidence that standards have declined in recent years, i.e., the post-war period.

For outbreaks in which the implicated food was investigated bacteriologically, and in those cases where epidemiological information regarding the suspected food was recorded, the main sources of infection were eggs and food containing eggs (33%), milk and milk products (22%), and minced beef (18%). Although this supports previous findings that eggs and egg products are the commonest vehicles of *S. Enteritidis* transmission [2,4], the large proportion of cases involving raw milk (22%) suggests that cattle from rural areas may represent an important source of infection. However, because of limited resources, food samples were examined from only five (19%) *S. Enteritidis* outbreaks during this 3-year period. Although the surveillance system for the sources of *S. Enteritidis* infections in this region is inadequate, this is also a problem in many other countries [1,2]. Medical staff in general practice should be strongly encouraged to report suspected cases to local public health authorities, and consistent criteria should be developed for their investigation.

## REFERENCES

1. Lindqvist R, Andersson Y, Lindbäck J *et al.* A one-year study of foodborne illnesses in the municipality of Uppsala, Sweden. *Emerg Infect Dis* 2001; **7**: 588–592.
2. Caffer MI, Eiguer T. *Salmonella enteritidis* in Argentina. *Int J Food Microbiol* 1994; **21**: 15–19.
3. Kelly MT, Brenner DJ, Farmer. JJ. Enterobacteriaceae. In: Lennette EH, Balows A, Hausler WJ, Shadomy HJ, eds. *Manual of clinical microbiology*. Washington DC: American Society for Microbiology, 1985; 263–277.
4. Duguid JP, North RAE. Eggs and salmonella food-poisoning: an evaluation. *J Med Microbiol* 1991; **34**: 65–72.
5. Centers for Disease Control and Prevention. *Update on Salmonella serotype Enteritidis infections*. Atlanta: Centers for Disease Control and Prevention, 1999.

## RESEARCH NOTE

### Increasing incidence of resistance to nalidixic acid in shigellas from humans in England and Wales: implications for therapy

T. Cheasty, M. Day and E. J. Threlfall

Health Protection Agency, Specialist and Reference Microbiology Division, Laboratory of Enteric Pathogens, London, UK

## ABSTRACT

Among shigellas isolated from patients in England and Wales in 2002, 10% of subgroups A, B and C, and 13% of subgroup D (*Shigella sonnei*), were resistant to nalidixic acid. As a consequence, should antimicrobial therapy be indicated, the efficacy of nalidixic acid as the preferred treatment for children with bacillary dysentery has been jeopardised.

**Keywords** Dysentery, nalidixic acid, quinolones, resistance, *Shigella*, therapy

**Original Submission:** 8 March 2004; **Revised Submission:** 20 May 2004; **Accepted:** 26 May 2004

*Clin Microbiol Infect* 2004; **10**: 1033–1035  
10.1111/j.1469-0691.2004.00982.x

The normal presentation of bacillary dysentery caused by *Shigella* isolates of subgroups A, B, C (*Shigella dysenteriae*, *S. flexneri*, *S. boydii*) and D (*S. sonnei*) is that of mild-to-moderate gastroenteritis. The disease is self-limiting, and the primary therapy is oral rehydration. However, symptoms can be severe in the very young, the very old, the malnourished and patients with other underlying diseases [1]. In such cases, administration of an effective antimicrobial agent should commence as soon as the clinical diagnosis is made. Ampicillin was the drug of choice until the mid-1980s [2], but this agent was compro-

Corresponding author and reprint requests: E. J. Threlfall, Health Protection Agency, Specialist and Reference Microbiology Division, Laboratory of Enteric Pathogens, 61 Colindale Avenue, London NW9 5HT, UK  
E-mail: john.threlfall@hpa.org.uk